

## Literaturauswahl zu Methoden zur Bestimmung der Körperzusammensetzung

(s. auch Abb. 34)

Keys, A. Brozek, J.: Body fat in adult man. *Physiological Reviews*, 33(3):245-25(1953) Lit. 7588

Brozek, J., Henschel, A. (eds.): *Techniques for measuring body composition*. Nat. Acad. Sci., Nat. Res. Council, Washington, D.C., 1961

Lantsche, W., Seige, K.: Über Möglichkeiten einer Bestimmung des Fettanteils am Körpergewicht. *Z. ges. inn. Med. Grenzgeb.*, 29:953-959(1974) Lit. 1356

Roche, A. F.: Some aspects of the criterion methods for the measurement of body composition. *Human Biology*, 59(2):209-220(1987) Lit. 19.919

Sheng, H.-P., Huggins, R. A.: A review of body composition studies with emphasis on total body water and fat. *Amer. J. clin. Nutr.*, 32: 630647(1979) Lit. 8568

Katch, F. I., McArdle, W. D.: Validity of body composition prediction equations for college men and women. *Amer. J. clin. Nutr.*, 28:105-109(1975) Lit. 8558

Smith, D. P., Boyce, R. W.: Prediction of body density and lean body weight in female 25 to 37 years old. *Amer. J. clin. Nutr.*, 30:560-564(1977) Lit. 8560

Garrow, J. S. et al.: A new method for measuring the body density of obese adults. *Brit. J. Nutr.*, 42:173-183(1979) (densitometry)

Katch, F. I., Katch, V. L.: Computer technology to evaluate body composition, nutrition and exercise. *Prev. Medicine*, 12:619-31(1983) Lit. 11.431 (densitometry)

Johnston, F. E., Paolone, A. N., Taylor, H. L., Schell, L. M.: The relationship of body fat weight, determined densitometrically to relative weight and triceps skinfolds in American youths, 12-17 years of age. *Amer. J. Physic. Anthropol.*, 57(1):1-6(1982) Lit. 12.449

Stock, M., Rothwell, N.: *Obesity and leanness – Basic Approach.*, K. Libbey, London, Paris, 82 p. (1982) (selbst separat)

Brozek, J., Grande, F., Anderson, J. T. Keys, A.: Densitometric analysis of body composition: revision of some quantitative assumptions. *Ann. N. Y. Acad. Sci.*, 110:113-140(1963)

Jones, F. R. M.: A body volumeter to measure human body density. *J. Physiol*, 222:5-7(1972)

Ward, A. et al.: A comparison of body fat determined by underwater weighing and volume displacement. *Amer. J. Physiol.*, 234(1):E49-E96(1978) CC 21(10)140

Opliger, R. A. et al.: Reliability of hydrostatic weighing and skinfold measurements of body composition using a generalizability study. *Human Biology*, 59(1):77-96(1987) CC 30(15)168

Oppliger, R. A., Looney, M. A., Tipton, Ch. M.: Reliability of hydrostatic weighing and skinfold measurements of body composition using a generalizability study. *Human Biology* 59(1):77-96 Lit. 19.941

Weltmann, A., Janney, C., Huber, R., Rians, C. B., Katchl, F. I.: Comparison of hydrostatic weighing at residual volume and total lung capacity in pre-pubertal males. *Human Biology*, 59(1):51-57(1987) Lit. 19.943

Mukherjee, D., Roclhe, A. F.: The estimation of percent body fat, body density and total body fat by maximum  $r^2$  regression equations. *Human Biology*, 56(1):79-109(1984) Lit. 19.868

Durnin, J. V. G. A., Rahaham, M. M.: The assessment of the amount of fat in human body from measurements of skinfold thickness. *Brit. J. Nutr.*, 21:681-689(1967)

Durnin, J. V. G. A., Womersley, J.: Body at assessed from total body density and its estimation from skinfold thickness: measurements on 481 men and women aged from 16 to 72 years. *Brit. J. Nutr.*, 32:77-97(1974) Lit. 8567

Committee on Nutrition: Measurement of skinfold thickness in childhood. *Pediatrics*, 42(3):538-543(1968) Lit. 225

Hermansen. L., Döberin, W. V.: Bodyfat and skinfold measurements. *Scand. J. clin. Lab. Invest.*, 27:315-319(1971)

Möhr, M.: Zur Definition der Fettsucht. *Ernährungsforschung*, 27(1):5-7(1982) Lit. 8780

Garrow, J. s.: Indices of adiosity. *Nutr. Abstr. Rev.*, 53(8):697-708(1983) Lit. 10.502

Sheng, H.-P., Huggins, R. D.: A review of body composition studies with emphasis on total body water and fat. *Amer. J. clin. Nutr.*, 32(3):630-647(1979) Lit. 8568

Lohman, T. G.: Skinfolde and body density and their relation to body fatness. A review. *Human Biology*, 53(2):181-226(1981) Lit. 8216

Martin, A. D., Ross, W. D., Drinkwater, D. T., Clarys, J. P.: Prediction of body fat by skinfold caliper : assumption and cadaver evidence. *Int. J. Obesity*, 9(Suppl. 1):31-39(1985) Lit. 14.531

Watson, P. E., Watson, I. D., Batt, R. D.: Total body water volumes for adult males and females estimated from simple anthropometric measurements. *Amer. J. clin. Nutr.*, 33:27-39(1980) Lit. 6470

Satwanti, Km., Bharadwaj, H., Singh, I. P.: Relationship of body density to body measurements in young Funjabi women: applicability of body composition prediction

equations developed for women of European descent. *Human Biology*, 49(2).203-213(1977) Lit. 3342

Andrzejewski, L., Hecking, E., Berres, M.: Messung der Tricaps-Hautfaltendicke. Aussagefähigkeit und Methodik. *Akt. Ernährungsw.*, 23(1):1-11(1984) Lit. 11.395

Pallier, E., Max, J. P., Burlet, C., Debry, G.: Human and animal fat cell size determination using an image analyzing computer. *Amer. J. clin. Nutr.*, 41:818-820(1985) Lit. 13.304

Cassady, C.: Bromide space studies in infants of low birth weight. *Pediat. Res.*, 4:14-24(1970) NAR 40:5090

Krzywicki, H. J., et al.: A comparison of methods for estimating human body composition. *Amer. J. clin. Nutr.*, 27:1380-1385(1974) (Densitometrie, Verdünnung, 40K, D20)

Boutton, T. W., Trowbridge, F. L., Nelson, M. M., Wills, C. A., Smith, E. O. Br., Lopez de Romana, G., Madrid, s., Marks, J. M., Klein, P. D.: Body composition of Peruvian children with short stature and high weight-for-height. I. Total body-water measurements and their prediction from anthropometric values. *Amer. J. clin. Nutr.* 45:513-525(1987) Lit. 17.963

Schoeller, D. A., Santen, E. van, Peterson, D. W., Dietz, W., Jaspán, J., Klein, P. D.: Total body water measurements in humans with  $^{18}\text{O}$  and  $^2\text{H}$  labeled water. *Amer. J. clin. Nutr.* 33:2686-2693(1980)

Trowbridge, F. L., Graham, G. G., Wong, W. W. et al.: Bodywater measurements in premature and older infants using  $\text{H}_2$   $^{18}\text{O}$  isotopic determination. *Ped. Res.*, 18:524-527(1984)

Fuchs, R. J., Theis, C.-F., Lancaster, M. C.: A monogram to predict lean body mass. *Amer. J. clin. Nutr.* 31:673-678(1978) Lit. 8563 (mit  $\text{T}_2\text{O}$ -Verdünnungstechnik)

Prentice, T. C., Siri, W. E., Berline, N. I., Hyde, G. M., Parsons, R. J., Joiner, E. E., Lawrence, J. H.: Studies of total body water with tritium. *J. clin. Invest.*, 31:412 (1952)

Szeluga, D. J. et al.: Nutritional assessment by isotope dilution analysis of body composition. *Amer. J. clin. Nutr.* 40: 847-854(1984)

Leonhardt, W., Hanefeld, M., Julius, U., Schulze, J., Fischer, S., Weck, m.; Haller, H. (Dresden): Predictive value of the index of desirable body weight for total body fat mass as measured by dilution of tritiated water – problems and limitations. *Int. J. Obesity*, 11:221-228(1987) Lit. 19.190

Lukaski, H. C., Johnson, P. E.: A simple inexpensive method of determining total body water using a tracer dose of  $\text{d}_2\text{O}$  and infrared absorption of biological fluid. *Amer. J. clin. Nutr.*, 41:363-370(1985)

Wong, W. W., Cochran, W. J., Klish, W. J., Smith, E. O. B., Lee, L. S., Klein, P. D.: In vivo isotope-fractionation factors and the measurements of deuterium and oxygen-

18-dilution spaces from plasma, saliva, respiratory water vapour and carbon dioxide. Amer. J. clin. Nutr. 47(1):1-6(1988) Lit. 20.733

Hanson, K. B., Garcia, P. A., Hotchkiss, D. K.: Estimation of body composition from anthropometric and physiological measurements of obese college women. Nutr. Res., 5:1055-1066(1985) Lit. 15.676 (Helium dilution)

Morgan, D. B., Burkinshaw, L.: Estimation of non-fat body tissues from measurements of skinfold thickness, total body potassium and total body nitrogen. Clin. Sci., 65(4):407-414(1983) NAR 54:2108

Webster, J. D., Hesp, R., Garrow, J. S.: The composition of excess weight in obese women estimated by body density, total body water and total body potassium. Human Nutr. Clin. Nutr., 38C:299-306(1984) Lit. 12.015

Lloyd, R. D., Mays, Ch. W.: A model for human body composition by total body counting. Human Biology, 59(1):7-30(1987) Lit. 19.920

Womersley, J. et al.: A comparison of the fat-free mass in young adults estimated by anthropometry, body density and total body potassium content. Clin. Sci. 43(3):469-475(1982) NAR 7262(1973)

Maaser, R., Droese, W., Würtenberger, H.: die Beurteilung des Ernährungszustandes. Ein Vergleich zwischen Ultraschall- und Calipermessung. KlinWschr., 50:923-926(1972) Lit. 448

Whittingham, F. D. G. V.: Measurement of tissue thickness by ultrasound. Aerospace Med., 33:1121(1962)

Sanchez, C. L., Jacobson, H. N.: Anthropometry measurements, a new type. Amer. J. clin. Nutr. 31(7):1116-1117(1978)

Popp, R. L., Macovski, A.: Ultrasonic diagnostic instruments. Science, 210:268-273(1980)

Paerisch, I., Paerisch, M.: Messung der Schichtdicke des menschlichen Unterhautfettgewebes mit dem Ultraschall-Impulsechoverfahren. Arch. ges. Physiol., 276:437(1983)

Bugyi, B.: Ultraschall-Echolot-Bestimmung des Unterhautfettgewebes und des Körperfettes. Z. Ernährungsw., 15:59-63(1976)

Hansen, W. E., Kehrer, H.: Assessment of cutaneous fat and body fat by ultrasound. Klin. Wschft., 65(9):407-410(1987) CC 30(21)173

Katch, F. I.: Individual differences of ultrasound assessment of subcutaneous fat: effects of body position. Human Biology, 55(4):789-796(1983) CC27(17)117

Blata, P. J., Ward, M. W. M., Tomkins, A. M. : Ultrasound for measurement of subcutaneous fat. Lancet, i(8218):504-505(1981)

Fanelli, M. T., Kuczmarski, R. J.: Ultrasound as an approach to assessing body composition. *Amer. J. clin. Nutr.*, 39:703-709(1984) Lit. 11.832

Kuczmarski, R. J., Fanelli, M. T., Koch, G. K.: Ultrasonic assessment of body composition in obese adults: overcoming the limitations of the skinfold caliper. *Amer. J. Clin. Nutr.*, 45:717-724(1987) Lit. 17.879

Conway, J., Norris, K., Bodwell, C. E.: A new approach for the determination of body composition in humans, infrared interactance. *Int. J. Obesity*, 9(Suppl. 2): A39(1985) Lit. 14.530 und *Amer. J. clin. Nutr.* 40(6):1123-1130(1984) Lit. 12.720

Weits, T., Beck, E. J. van der, Wedel, M.: Comparison of ultrasound and skinfold caliper measurements of subcutaneous fat tissue. *Inter. J. Obesity*, 10:161-168(1986) Lit. 17.602 und 16.010

Chumlea, W. C., Roche, A. F.: Ultrasonic and skinfold caliper measures of subcutaneous adipose tissue thickness in elderly men and women. *Amer. J. Phys. Anthropol.*, 71(3):351-358(1986)

Jones, P. R. M. et al.: Ultrasonic measurements of subcutaneous adipose tissue thickness in men. *Amer. J. physic. Athrop.* 71(3):359-364(1986) CC 29(50) 155

Garrow, J. S.: New approaches to body composition. *Amer. J. clin. Nutr.*, 35(5. Suppl.):1152-1158(1982) selbst separat

Harrison, G. G., Itallie, T. B. van: Estimation of body composition: a new approach based on electromagnetic principles. *Amer. J. clin. Nutr.* 35:1176-1179(1982)

Presta, E., Wang, J., Harrison, G. G., Björntolrp, P., Harker, W. H., Itallie, T. B. van: Measurement of total body electrical conductivity: a new method for estimation of body composition. *Amer. J. clin. Nutr.*, 37:735-739(1983) und Vol. 35:824Abs.(1982) Lit. 9073

Presta, E., Segal, K. R., Gutin, B., Harrison, G. G., Itallie, T. B. van: Comparison in man of total body electrical conductivity and lean body mass derived from body density: validation of a new body composition method. *Metabolism*, 35:524-527(1983)

Segal, K. R., Gutin, B., Presa, E., Wang, J., Itallie, T. B. van: Estimation of human body composition by electrical impedance methods: a comparative study. *J. Appl. Physiol.* 58:1565-1571(1985)

Chochran, W. J. et al.: Total body electrical conductivity used to determine body composition in infants. *Ped. Res.* 20(6):561-564(1986)

Lucasi, H. C., Johnson, P. E., Bolonchuk, W. W., Lykken, G. I.: Assessment of fat-free mass using bioelectrical impedance measurements of the human body. *Amer. J. clin. Nutr.*, 41:810-817(1985) Lit. 13.363

Pasco, J. A., Rutishauser, I. H. E.: Body fat estimated from anthropometric and electric impedance measurements. *Human Nutr. Clin. Nutr.*, 39C:365-369(1985) Lit. 14.672

Cohn, s. H.: How valid are bioelectric impedance measurements in body composition studies? Amer. J. clin. Nutr., 42:889-890(1985) Lit. 14.750

Kushner, R. F., Schoeller, D. A.: Estimation of total body water by bioelectrical impedance analysis. Amer. J. clin. Nutr., 44:417-424(1986) Lit. 17.083

Solomons, N. W.: How valid are bioelectric impedane measurements in body composition study? Amer. J. clin. Nutr., 44:306-307(1986) Lit. 17.084

Mazess, R. B.: Accuracy of TOBEC method questioned. Amer. J. clin. Nutr., 39(1):157-159(1984) Lit. 11.158

Loan, M. van, Maycline, P.: A new TOBEC instrument and procedure for the assessment of body composition: use of Fourir coefficients to predict lean body mass and total body water. Amer. J. clin. Nutr., 45: 131-137(1987) Lit. 17.409

Loan, M. D. van, Belko, A. Z., Mayclin, P. L., Barbieri, T. F.: Use of total-body electrical conductivity for monitoring body composition changes during weight reeducation. Amer. J. clin. Nutr., 46: 5-8(1987) Lit. 18.713

Loan, M. D. van, Segal, K. R., Bracco, E. F., Maycline, P., Itallie, T. B. van. TOBEC methodology for body composition assessment: a gross validation study. Amer. J. clin. Nutr. 46: 9-12(1987)

The bioelectrical estimation of body composition. Human Biology, 59(No. 2, April 1987 (ganzes Heft zu diesem Thema, S. 209-336) CC30(18) 135(1987) Lit. 17.880

Daraus z.B. Malina, R. M.: Bioelectric methods for estimating body composition: an overview and discussion, p. 329-335, Lit. 19.912

p. 319-327: Klish, W. J., Cochran, W. J., Fiorotto, M. L., Wong, W. W., Klein, P. D.: The bioelectrical measurements of body composition during infancy. Lit. 19.913

p. 281-298: Hodgdon, J. A., Fitzgerald, P. I.: Validity of impedance predictions at various levels of fatness.

p. 257-269: Cumlea, Wm. C., Roche, A. F., Guo, Shl, Woynarowska, B.: The influence of physiologic variables and oral contraceptives on bioelectric impedance.

p. 221-233: Guo, Sh., Roche, A. F., Chumlea, Wm. C., Miles, D. S., Fohlmann, R. L.: Body composition prediction from bioelectric impedance. Lit. 19.917

Segal, K. R., Loan, M. van, Fitzgerald, P. I., Hodgdon, J. A., Iallie, T., B. van: Lean body mass estimation by bioelectrical impedance analysis: a four-site cross-validation study. Amer. J. clin. Nutr. 47(1):7-14(1988) Lit. 20.734

Cunningham, J J.: New approaches to the noninvasive assessment of body composition: bioelectrical impedance analysis and total body electrical conductivity. Nutr. Internat., 3(1):6-11(1987) CC 30(12) 80 Lit. 20.261

Diaz, E., Villar, J., Immink, M., Gonzalez, T.: Bioimpedance analysis: is it a satisfactory method for the assessment of body composition. Proc. Nlutr. Soc., 47:57A(1988) Lit. 20.516

Kushner, R. F., Haas, A.: Estimation of lean body mass by bioimpedance analysis compared to skinfold anthropometry. *Amer. J. clin. Nutr.* 45: 830 abs (1987)

Lubanski, R. E., Kurzer, M., Meguid, M. M.: Measurement of body composition using bioelectrical impedance analysis. *Amer. J. clin. Nutr.*, 45:830abs(1987) Lit. 20.539

Schell, B., Gross, R.: The reliability of bioelectrical impedance measurements in the assessment of body composition in healthy adults. *Nutr. Rep. Intern.*, 36(2):449-459(1987 Lit. 19.930

Katch, I., Solomon, T. T., Shayevitz, M., Shayevitz, B.: Validity of bioelectrical impedance to estimate body composition in cardiac and pulmonary patients. *Amer. J. clin. Nutr.*, 43:972-973(1986) Lit. 15.878

Katch, F. I., Katch, V. L.: Computer technology to evaluate body composition, nutrition and exercise. *Preventive Medicine*, 12:619-631(1983) Lit. 11.431

Tanner, J. M. et al.: Radiographically determined width of bone, muscle and fat in the upper arm and calf from age 3-18 years. *Ann. Human Biology*, 8(6):495-518(1981) CC 25(2) 150

Katch, F. I., Behnke, A. R.: A X-ray assessment of percent body fat in man and women. *Med. Sci. Sport Exercise*, 16(3):316 ff(1984) CC 27(29) 147

Tokunaga, K. et al.: A novel technique for the determination of body fat by computed tomography. *Int. J. Obesity*, 7:437-445(1983)

Kvist, H., Sjöström, L., Tylén, U.: Adipose tissue volume determination in women by computed tomography: technical considerations. *Int. J. Obesity*, 10:53-67(1986) Lit. 15.203

Seidell, J. C., Oosterlee, A., Thijssen, M. A. O., Burema, J., Deurenberg, P., Hautvast, J. G. A. J., Ruijs, J. H. J.: Assessment of intra-abdominal and subcutaneous abdominal fat: relation between anthropometry and computed tomography. *Amer. J. clin. Nutr.*, 45:7-13(1987) Lit. 17.408

Ashwell, M., Cole, T. J., Dixon, A. K.: Obesity: new insight into anthropometric classification of fat distribution shown by computed tomography. *Brit. Med. J.*, 290:1692-1694(1985)

Dixon, A. K.: Abdominal fat assessed by computed tomography: sex difference in distribution. *Clin. Radiology*, 14:189-191(1983)

Borkan, G. A., Gerzof, S. G., Robbins, A. H., Hulst, D. E., Silbert, C.K.: Assessment of abdominal fat content by computed tomography. *Amer. J. clin. Nutr.* 36:172-177(1982)

Borkan G. A., Hulst, D. E., Gerzof, S. G., Robbins, A. H., Silbert, C. K.: Age changes in body composition revealed by computed tomography. *J. Gerontology*, 38:673-677(1983)

Grauer, W. O., Moss, A. A., Cann, C. E., Goldberg, H. I.: Quantification of body fat distribution in the abdomen using computed tomography. *Amer. J. clin. Nutr.*, 39:631-637(1984)

Tlapák, P., Otáhal, S., Hrabé, J., Parisková, J.: Estimation of muscle, fat, bone and intestine in pelvic sections of individuals with different body mass indices. *Human Nutr. Clin. Nutr.*, 40C:393-395(1986) Lit. 16.714 (Tomography)

Huang, H. K., Wu, S. C.: The evaluation of mass densities of the human body in vivo from CT from CT scans. *Comput. Biol. Med.*, 6:337-343(1976)

Borkan, G. A. et al.: Relationships between computed tomography tissue areas, thickness and total body composition. *Ann. Human Biology*, 10(6):537-546(1983) CC 27(2) 128

Enzi, G. et al.: Subcutaneous and visceral fat distribution according to sex, age and overweight evaluated by computer tomography. *Amer. J. clin. Nutr.*, 44:739-746(1986)

Buckley, D. C., Kudsk, K. A., Rose, B. S., Fatzinger, P., Koetting, C. A.a, Schlatter, M.: Anthropometric and computerized komographic measurements of lower extremity lean body mass. *J. Amer. Diet. Assoc.*, 87(2):196-199(1987) Lit. 20.181

Mazess, R. B., Peppler, W. W., Gibbons, M.: Total body composition by dual-photon ( $^{153}\text{Gd}$ ) absorptiometry. *Amer. J. clin. Nutr.*, 40:834-839(1984) Lit. 12.317

Lusaki, H. C. et al.: A comparison of methods of assessment of body composition including neutron activation analysis of total body nitrogen. *Metabolism*, 30(8):777-782(19881) CC 24(37)122

Cohn, S. H. et al.: Improved models for determination of body fat by in vivo neutron activation. *Amer. J. clin. Nutr.*, 40(2):255-259(1984)

Almond, D. J. et al.: Measurement of short-term changes in the fat content of the vody: a comparison of three methods in patients receiving intravenous nutrition. *Brit. J. Nutr.*, 52:215-225(1984) (neutron activation)

Harrison, G. G., Itallie, T. B. van: Estimation of body composition – a new approach based on electromagnetic principles. *Amer. J. clin. Nutr.*, 35 (5. Suppl.): 1176-1179(1982 selbst separat

Khaleb, M. A., Lukaski, H. C., Watkins, C. L.: Determination of total body water by deuterium NMR. *Amer. J. clin. Nutr.*, 45:1-6(1987)

Rebouche, Ch. J. et al.: Evaluation of nuclear magnetic responance spectroscopy for determination of deuterium abundance in body fluids: application to measurement of total body water in human infants. *Amer. J. clin. Nutr.*. 45:373-380(1987)

Ulin, K., Meydani, M., Zamenhof, R. G., Blumberg, J. B.: Photon activation analysis as a new technique for body composition studies. *Amer. J. clin. Nutr.*, 963-972(1986) Lit. 16.850



Wang, J., McKeon, E., Heymsfield, S., Kral, J., Pierson, R. N. jr.: A new framework for body composition studies: potassium, hydration, and density in the lean body mass measured by dual photon absorptiometry. *Amer. J. clin. Nutr.* 45:833abs(1987)  
Lit. 20.545

Cohn, S. H., Brennan, B. L., Yasumura, S., Vartsky, D., Vaswami, A. N., Ellis, K. J.: Evaluation of body composition and nitrogen content as determined by total body neutron activation. *Amer. J. clin. Nutr.*, 38:52-58(1983)